

9. (Previously presented) The method of claim 6, wherein the monoclonal antibody is a chimeric monoclonal antibody containing constant regions and variable regions from different species.
10. (Previously presented) The method of claim 6, wherein the monoclonal antibody is a humanized monoclonal antibody.
11. (Canceled)
12. (Currently amended) ~~The method of claim 1,~~ A method for preventing a T cell mediated autoimmune response associated with type I diabetes comprising administering to a subject in need of such treatment a prophylactically effective amount of wherein the gp39-antagonist is an anti-gp39 antibody or a gp39-binding fragment thereof, comprising variable regions of monoclonal antibody 24-31 or monoclonal antibody 89-76, or of an antibody having the gp39 binding characteristics thereof.
13. (Previously presented) The method of claim 12, wherein the gp39 antagonist is a gp39-binding antibody fragment comprising variable regions of monoclonal antibody 24-31.
14. (Previously presented) The method of claim 13, wherein the gp39-binding antibody fragment is a Fab or F(ab')₂ fragment comprising variable regions of monoclonal antibody 24-31.
15. (Previously presented) The method of claim 12, wherein the gp39 antagonist is a gp39-binding antibody fragment comprising variable regions of monoclonal antibody 89-76.
16. (Previously presented) The method of claim 15, wherein the gp39-binding antibody fragment is a Fab or F(ab')₂ fragment comprising variable regions of monoclonal antibody 89-76.

17. (Previously presented) The method of claim 9, wherein the chimeric monoclonal anti-gp39 antibody comprises variable regions of monoclonal antibody 24-31.
18. (Previously presented) The method of claim 9, wherein the chimeric monoclonal anti-gp39 antibody comprises variable regions of monoclonal antibody 89-76.
19. (Previously presented) The method of claim 10, wherein the humanized monoclonal anti-gp39 antibody comprises variable regions of monoclonal antibody 24-31.
20. (Previously presented) The method of claim 10, wherein the humanized monoclonal anti-gp39 antibody comprises variable regions of monoclonal antibody 89-76.
21. (New) A method of preventing a T cell mediated autoimmune response associated with type I diabetes comprising the method of claim 1.